

# **N Bases / Nucleosides / Nucleotides / Nucleic Acid Structures (Review)**

## **Goals for this review unit:**

- 1. Recognize the common building blocks of nucleic acids: names / 1-letter abbrev.**
- 2. Nomenclature for nucleosides and nucleotides (structure of ATP)**
- 3. Primary structures of RNA and DNA**
- 4. Conformations in DNAs**
- 5. Characteristics of B-DNA, A-DNA and Z-DNA**
- 6. Denaturation of DNA**
- 7. Features of RNA / Functions of RNA**
- 8. DNA Sequencing (Maxam – Gilbert vs. Sanger Dideoxy)**

## iClicker Question #1

The approximate charge at pH 1 of the oligopeptide “H – A – P – P – Y – D – A – Y – S” is \_\_\_\_.

Answers: A) +3 B) +2 C) +1 D) -1 E) -2

## iClicker Question #2


**The world at 1,000,000 X ( 1 nm → 1 mm )**

At this magnification, double stranded DNA would be approximately \_\_\_\_\_ in width.

Answers: A) 1 mm B) 1 m C) 2 mm D) 20 m E) 5 mm

# The Birth of Molecular Biology: DNA Structure

inside, linked together by hydrogen bonds. This structure as described is rather ill-defined, and for this reason we shall not comment on it.



This figure is purely diagrammatic. The two ribbons symbolize the two phosphate-sugar chains, and the horizontal rods the pairs of bases holding the chains together. The vertical line marks the fibre axis

We wish to put forward a radically different structure for the salt of deoxyribose nucleic acid. This structure has two helical chains each coiled round the same axis (see diagram). We have made the usual chemical assumptions, namely, that each chain consists of phosphate di-ester groups joining  $\beta$ -D-deoxy-ribofuranose residues with 3',5' linkages. The two chains (but not their bases) are related by a dyad perpendicular to the fibre axis. Both chains follow right-handed helices, but owing to the dyad the sequences of the atoms in the two chains run in opposite directions. Each chain loosely resembles Furberg's<sup>2</sup> model No. 1; that is, the bases are on the inside of the helix and the phosphates on the outside. The configuration of the sugar and the atoms near it is close to Furberg's 'standard configuration', the sugar being roughly perpendicular to the attached base. There

*Nature* – 1953



*Nature* – 2001





## The Nobel Prize in Physiology or Medicine 1962

"for their discoveries concerning the molecular structure of nucleic acids and its significance for information transfer in living material"



**Francis Harry Compton Crick**

🕒 1/3 of the prize

United Kingdom

MRC Laboratory of Molecular Biology  
Cambridge, United Kingdom

b. 1916  
d. 2004



**James Dewey Watson**

🕒 1/3 of the prize

USA

Harvard University  
Cambridge, MA, USA

b. 1928



**Maurice Hugh Frederick Wilkins**

🕒 1/3 of the prize

United Kingdom and New Zealand

London University  
London, United Kingdom

b. 1916  
(in Pongaroa, New Zealand)  
d. 2004





Left to right: Maurice Wilkins, John Steinbeck, John Kendrew, Max Perutz, Francis Crick and Jim Watson after the Nobel Ceremony in Stockholm in December 1962.


## Pyrimidines



## Purines




**Deoxyribose (in DNA)**




**Ribose (in RNA)**

**(a) Nucleotide components**







**(b) Nucleotide**



**(c) Polynucleotide**

iClicker Question #3 : What is the correct name for #2?



A) Guanosine


B) Thymine

C) Cytidine MP


D) deoxyThymine MP

E) dexoyThymidine MP







**dAdenosine MP**



**dThymidine MP**




**dGuanosine MP**




**dCytidine MP**

# DNA primary structure







# Rotational angles of phosphodiester chain




	B-DNA	A-DNA	Z-DNA C	Z-DNA G
$\alpha(\omega)$	-41	-90	138	100
$\beta(\phi)$	136	211	-94	-108
$\gamma(\psi)$	38	47	80	-70
$\delta(\psi')$	139	83	48	-130
$\epsilon(\phi')$	-133	-185	180	-140
$\zeta(\omega')$	-57	-45	-170	56
$\chi$	78	27	20	-100







guanosine-anti




guanosine-syn



C3' endo




C2' exo






C3' endo-C2' exo


# Sugar pucker in DNA



$\beta$ -D-2-Deoxyribose




*$C3'$  endo*




*$C2'$  endo*

# Sugar pucker in DNA



5.9 Å

*C3' endo*




7.0 Å


*C2' endo*



# Double stranded DNA




- *Two single stranded DNA paired by Hydrogen bonds.*
- *Helical structure*




### homo purine

symmetric




asymmetric




### hetero purine

symmetric




asymmetric




### homo pyrimidine

symmetric




asymmetric




### hetero pyrimidine

symmetric only




### purine - pyrimidine


symmetric




asymmetric



# Watson-Crick base pairs




*G-C base pair*




*A-T base pair*




# A and B Double Helices




(a) B-DNA, end-on view




(c) A-DNA, end-on view



(b) B-DNA, side view




(d) A-DNA, side view





## Structure of double stranded DNA (B-DNA)



**B-DNA**

*Right handed helix*

*10.5 residue per turn*

*Helix pitch = 34Å*

*Base pair tilt-helical axis = -6°*

*Diameter = 20Å*

*Sugar pucker*

*dA, dT, dC, dA: C2' endo*

*Glycosidic bond*

*dA, dT, dC, dA: anti*

*Minor grove show base diversity*

## A-DNA vs. B. DNA

### A - DNA

*11 residue per turn*

*Helix pitch = 28Å*

*Base pair tilt = 20°*


*Diameter = 23Å*

*Sugar pucker*


*C3' endo*

*Glycosidic bond*



*anti*




B form




A form



# Z-DNA vs. B-DNA



B-DNA



Z-DNA

## Z - DNA

*left handed helix*

*12 residue per turn*

*Helix pitch = 45Å*

*Base pair tilt = 7°*

*Diameter = 18Å*

*Sugar pucker*

*dA, dT, dC: C2' endo*

*dG: C3' endo*

*Glycosidic bond*

*dA, dT, dC: anti*


*dG: syn*




# Denatuted DNA:

Heat denaturation of DNA is called "melting," The purine and pyrimidine bases exhibit very strong p-p\* transitions around 260 nm. *E. coli* DNA absorption is only about 60% of that predicted from the weighted average spectrum based on its composition, this loss of intensity is called **hypochromism**. Since the absorbance goes up as DNA "unwinds", it can be used to monitor the unstacking of DNA.


Campbell, Biochemistry, 3/e  
Text Figure 07.13




Harcourt Brace & Company





# RNA primary structure



# Transfer-RNA



*Nonlocal  
basepairs*





Ribosome

KICHKA@12.09

Ada Yonai


Nobel Prize



Head Full of Ribosomes









# The Proto-Ribosome: An Ancient Nano-machine for Peptide Bond Formation

Chen Davidovich,<sup>[a]</sup> Matthew Belousoff,<sup>[a]</sup> Itai Wekselman,<sup>[a]</sup> Tal Shapira,<sup>[a]</sup> Miri Krupkin,<sup>[a]</sup> Ella Zimmerman,<sup>[a]</sup> Anat Bashan,<sup>\*[a]</sup> and Ada Yonath<sup>[a]</sup>



**Figure 1.** Top: the symmetrical region within the large ribosomal subunit. Bottom: Schematic representation of "pocket-like" proto-ribosome formation from a RNA precursor, showing simple catalytic peptidyl transferase activity.

# Genetic information



... *G T A C T G A A C G C A G G T*...


*Genetic code*

*Human being: ~ 3,000,000,000 base-pairs*


*~ 30,000 – 40,000 Genes*

*(Public Human Genome Project and Celera Genomics)*

# Chromosome



*Human being has 24 chromosomes.*





# Sequencing DNA

Prior to the **mid-1970's no method** existed by which DNA could be directly sequenced. Knowledge about gene and genome organization was based upon studies of prokaryotic organisms and the primary means of obtaining DNA sequence was so-called **reverse genetics** in which the **amino acid sequence of the gene product** of interest is **back-translated** into a nucleotide sequence based upon the appropriate codons.

- **Maxam-Gilbert DNA Sequencing**
- **Sanger (didexoy) DNA Sequencing**



## The Nobel Prize in Chemistry 1958

"for his work on the structure of proteins, especially that of insulin"



**Frederick Sanger**

United Kingdom

University of Cambridge  
Cambridge, United Kingdom

b. 1918



## The Nobel Prize in Chemistry 1980

"for his fundamental studies of the biochemistry of nucleic acids, with particular regard to recombinant-DNA"

"for their contributions concerning the determination of base sequences in nucleic acids"



**Paul Berg**

🕒 1/2 of the prize

USA

Stanford University  
Stanford, CA, USA

b. 1926



**Walter Gilbert**

🕒 1/4 of the prize

USA

Harvard University,  
Biological Laboratories  
Cambridge, MA, USA

b. 1932



**Frederick Sanger**

🕒 1/4 of the prize

United Kingdom

MRC Laboratory of  
Molecular Biology  
Cambridge, United Kingdom

b. 1918

# Maxam-Gilbert DNA Sequencing




Figure 1. Chemical targets in the Maxam-Gilbert DNA sequencing strategy. Dimethylsulphate or hydrazine will attack the purine or pyrimidine rings respectively and piperidine will cleave the phosphate bond at the 3' carbon.

[http://www.idtdna.com/support/technical/TechnicalBulletinPDF/DNA\\_Sequencing.pdf](http://www.idtdna.com/support/technical/TechnicalBulletinPDF/DNA_Sequencing.pdf)



**IDT**Tutorial: DNA Sequencing

# Allan Maxam / Walter Gilbert DNA Sequencing

## Sequencing single-stranded DNA

Two-step catalytic process:

1) Break glycoside bond between the ribose sugar and the base /  
displace base

Purines react with dimethyl sulfate

Pyrimidines react with hydrazine

2) Piperidine catalyzes phosphodiester bond cleavage where base  
displaced

---

“G” - dimethyl sulfate and piperidine

“A + G” - dimethyl sulfate and piperidine in formic acid

“C” - hydrazine and piperidine in 1.5M NaCl

“C + T” - hydrazine and piperidine



5' \*pCpCpGpGpCpGpCpApGpApApGpCpGpGpCpApTpCpApGpCpApApA 3'




Figure 2. The Maxam-Gilbert manual sequencing scheme. The target DNA is radiolabeled and then split into the four chemical cleavage reactions. Each reaction is loaded onto a polyacrylamide gel and run. Finally, the gel is autoradiographed and base calling proceeds from bottom to top.

# Maxam-Gilbert DNA Sequencing




- 200-300 bases of DNA sequence every few days
- Use large amounts of radioactive material,  $^{35}\text{S}$  or  $^{32}\text{P}$
- Constantly pouring large, paper thin acrylamide gels
- **Hydrazine is a neurotoxin**

## *Early Benefits -*

*Discovery that the gene for ovalbumin in chicken and the gene encoding  $\beta$ -globin in rabbit contained **non-coding gaps** in the coding regions. These gaps were flanked by the same dinucleotides in the two genes; GT on the 5' end of the gaps and AG on the 3' end of the gaps. Soon, the terms **intron** and **exon** were added to the genetic lexicon to describe the coding and non-coding regions of eukaryotic genes (1977).*




# Fred Sanger (dideoxy) DNA Sequencing

Sanger knew that, whenever a dideoxynucleotide was incorporated into a polynucleotide, the chain would irreversibly stop, or terminate. Thus, the **incorporation of specific dideoxynucleotides** in vitro would result in **selective chain termination**.



ddNTP analog

# Sanger (dideoxy) DNA Sequencing






Consider the following nucleic acid sequencing gel experiment using the Sanger dideoxy sequencing method:

What is the expected sequence (5' -> 3') of the **original** DNA sample assuming the primer was labeled with a 5'-prime fluorescent label?

[ DNA polymerase I + 4 dNTPs + ddATP ddTTP ddCTP ddGTP ]  
**A T C G**

**iClicker Question #4:**

- A) GACTTGA
- B) CTGAACT
- C) AGTTCGA
- D) TCAAGTC
- E) None of the above



## Advantages of dideoxy DNA Sequencing

- **Elimination of dangerous chemicals (hydrazine)**
- **Greater efficiency**


**Taq polymerase makes DNA strands off of a template at rate of about 500 bases per minute**

**Chemical synthesis of a 25-mer oligonucleotide takes more than two hours.**

→ **High Throughput Methods (Human Genome Project)**


# Automated Fluorescence Sequencing

In **1986**, Leroy Hood and colleagues reported on a DNA sequencing method in which the **radioactive labels, autoradiography, and manual base calling** were all replaced by **fluorescent labels, laser induced fluorescence detection, and computerized base calling**.



**Figure 5. A.** Chemical structure of the four succinylfluorescein dyes developed at DuPont. **B.** Normalized fluorescence emission spectra for each of the four dyes following excitation at 488nm. Shifts in the spectra were achieved by changing the side groups R<sub>1</sub> and R<sub>2</sub>.

# Automated DNA sequencing






# Automated dye-terminator sequencing

4-fluorescently labelled dideoxy dye terminators

ddATP  
ddGTP  
ddCTP  
ddTTP

pool and load in a single well or capillary

- scan with laser + detector specific for each dye
- automated base calling
- very long reads (~ 1000 bases)/run



[DNA dideoxysequencing animation](#)



# National Center for Biotechnology Information

National Library of Medicine

National Institutes of Health

PubMed All Databases **BLAST** OMIM Books TaxBrowser Structure

Search All Databases for  Go

- SITE MAP
  - Alphabetical List
  - Resource Guide
- About NCBI
  - An introduction to NCBI
- GenBank
  - Sequence submission support and software
- Literature databases
  - PubMed, OMIM, Books, and PubMed Central
- Molecular databases
  - Sequences, structures, and taxonomy

## What does NCBI do?

Established in 1988 as a national resource for molecular biology information, NCBI creates public databases, conducts research in computational biology, develops software tools for analyzing genome data, and disseminates biomedical information - all for the better understanding of molecular processes affecting human health and disease. [More...](#)

## 100 Gigabases

GenBank and its collaborating databases, the European Molecular Biology Laboratory and the DHA Databank of Japan, have reached a milestone of 100 billion bases from over 165,000 organisms. See the [press release](#) or find more information on [GenBank](#).

## CCDS Database


## Hot Spots

- Assembly Archive
- Clusters of orthologous groups
- Coffee Break, Genes & Disease, NCBI Handbook
- Electronic PCR
- Entrez Home
- Entrez Tools
- Gene expression omnibus (GEO)
- Human genome resources
- Malaria genetics & genomics

## International sequence databases exceed 100 gigabases

In August 2005, the INSDC announced the DNA sequence database exceeded 100 gigabases. GenBank is proud of its contributions toward this milestone. We thank all the scientists who have worked through the submission process at GenBank and made their sequence data available to the world. See the related [press release](#).

**>100,000,000,000 bases**



**> 200,000 organisms!!**